

## Research Note

### Domiciliary cockroaches found in restaurants in five zones of Kuala Lumpur Federal Territory, peninsular Malaysia

Jeffery, J.<sup>1</sup>, Sulaiman, S.<sup>2</sup>, Oothuman, P.<sup>3</sup>, Vellayan, S.<sup>4</sup>, Zainol-Arifin, P.<sup>5</sup>, Paramaswaran, S.<sup>6</sup>, Razak, A.<sup>2</sup>, Muslimin, M.<sup>1</sup>, Kamil-Ali, O.B.<sup>1</sup>, Rohela, M.<sup>1</sup> and Abdul-Aziz, N.M.<sup>1\*</sup>

<sup>1</sup>Department of Parasitology, Faculty of Medicine, University of Malaya, 50605 Kuala Lumpur, Malaysia

<sup>2</sup>Department of Biomedical Science, Faculty of Allied Health Sciences, Universiti Kebangsaan Malaysia, Jalan Raja Muda Abdul Aziz, 50300 Kuala Lumpur, Malaysia

<sup>3</sup>Kulliyah of Medicine, International Islamic University, P.O. Box 141, 25710 Kuantan, Pahang, Malaysia

<sup>4</sup>Faculty of Medicine, Universiti Teknologi MARA, 40450 Shah Alam, Selangor, Malaysia

<sup>5</sup>Public Health Division, Federal Territory of Kuala Lumpur and Putrajaya Health Office, Jalan Cenderasari, 50590 Kuala Lumpur, Malaysia

<sup>6</sup>Epidemiology and Biostatistic Unit, Institute for Medical Research, Jalan Pahang, 50588 Kuala Lumpur, Malaysia

\*Corresponding author email: noisha@ummc.edu.my

Received 19 July 2011; received in revised form 18 August 2011; accepted 20 August 2011

**Abstract.** The following domiciliary cockroaches were collected from restaurants in five zones of Kuala Lumpur Federal Territory, Malaysia using 1L glass beaker traps baited with ground mouse-pellets: *Periplaneta americana* (Linnaeus) (n = 820), *Periplaneta brunnea* Burmeister (n = 46), *Blattella germanica* (Linnaeus) (n = 12504), *Supella longipalpa* (Fabricius) (n = 321), *Symptloce pallens* Stephens (n = 29) and *Neostylopyga rhombifolia* (Stoll) (n = 5). The following bacteria were isolated from 10 cockroach specimens: *Enterobacter cloacae*, *Klebsiella pneumoniae* ssp. *pneumoniae*, *Klebsiella pneumoniae* ssp. *rhinoscleromatis* and *Serratia liquefaciens* from 5 *B. germanica*; *Acinetobacter calcoaceticus* var. *anitratus*, *Citrobacter diversus/amalonaticus*, *Escherichia vulneris* and *K.p. pneumoniae* from 3 *P. brunnea*; and *Citrobacter freundii*, *Enterobacter agglomerans* 4, *Escherichia adecarboxylate*, *E. vulneris*, *K. p. pneumonia*, *K. p. rhinoscleromatis* and *Proteus vulgaris* from 2 *P. americana*.

Cockroaches are a primitive and highly successful form of insect life (Cochran, 1999). They have exploited a variety of habitats in the tropics and subtropics (Cornwell, 1968). About 35 species, termed as domiciliary pest species, have established varying degrees of association in human dwellings and premises, and in peninsular Malaysia, 13 species have been collected in such situations (Oothuman *et al.*, 1984; Yap *et al.*, 1991; Lee *et al.*, 1993; Vythilingam *et al.*, 1997; Jeffrey *et al.*, 1999, 2003a, 2003b; Lee & Lee, 2000; Sulaiman *et al.*, 2003).

Cockroaches have been collected from several places including Klang (Oothuman *et al.*, 1984), Penang (Yap *et al.*, 1991; Lee *et al.*, 1993) and Kuala Lumpur (Vythilingam *et al.*, 1997; Sulaiman *et al.*, 2003). However, the collection from Kuala Lumpur was confined to the Chow Kit area, Taman Ibu Kota and Setapak Jaya.

Here cockroaches from 5 out of 6 zones of Kuala Lumpur Federal Territory and isolation of bacteria from 10 cockroach specimens are reported.

Cockroaches were trapped from the kitchen and other vantage areas of restaurants in 5 out of 6 zones of the Kuala Lumpur Federal Territory. Two beaker-traps baited with ground mouse-pellets (Jeffery *et al.*, 1984) were set up at each selected restaurant in the following areas: Taman Tun Dr. Ismail and Bangsar (Zone Damansara); Taman Wahyu, Kepong Baru and Jalan Kok Lian (Zone Kepong); Jalan Genting Klang and Jalan Tun Abdul Razak/Jalan Semarak (Zone Setapak); Jalan Peel and Taman Maluri (Zone Cheras); and 4 1/4 mile Jalan Klang Lama, Jalan Mega Mendung, Jalan Tun Sambanthan and 6<sup>th</sup> mile Jalan Puchong (Zone Klang Lama).

Weekly, trapped cockroaches were transferred into plastic bags and brought back to the laboratory for recording and identification. Cockroaches, except *Symploce pallens*, were identified using descriptions and keys in Cochran (1999) and compared with reference specimens maintained in the department. *Symploce pallens*, reported earlier by Jeffery *et al.* (1997), was initially identified by the late Roth, L.M. of Massachusetts, USA.

Ten cockroach specimens (5 *Blattella germanica*, 3 *Periplaneta brunnea* and 2 *Periplaneta americana*) were processed for the presence of bacteria. Isolation and identification were done according to the methods mentioned in Oothuman *et al.* (1989). Guts of the cockroaches were dissected out, under sterile conditions, and emulsified in petri dishes containing sterile normal saline. A small amount of the emulsion was transferred into sterile peptone water and a loopful of the suspension was inoculated on to Oxoid blood agar base containing 5% blood and MacConkey medium. The plates were incubated aerobically for 24-28 h at 37°C. Bacterial identification was carried out according to Cowan & Steel (1970).

A total of 13725 specimens (3204 females, 4337 males and 6184 nymphs) of 6 species of cockroach were trapped (Table 1). The dominant species trapped was the German cockroach, *B. germanica* (Linnaeus) (n = 12504, 91.10%); followed by

the American cockroach, *P. americana* (Linnaeus) (n = 820, 5.97%); the brown-banded cockroach, *Supella longipalpa* (Fabricius) (n = 321, 2.34%); the large brown cockroach, *P. brunnea* (Burmeister) (n = 46, 0.34%); the smooth cockroach, *S. pallens* (Stephens) (n = 29, 0.21%); and the harlequin cockroach, *Neostylopyga rhombifolia* (Stoll) (n = 5, 0.04%). *Blattella germanica*, *P. americana*, *P. brunnea* and *S. longipalpa* were trapped from all 5 zones; whereas *N. rhombifolia* and *S. pallens* were not trapped from the Setapak and Kepong zones, respectively.

*Blattella germanica* is prevalent in food outlets in peninsular Malaysia (Lee, 1998). In this study, it was trapped from all the zones. It was not reported from residential premises in earlier studies (Jeffery *et al.*, 1982; Oothuman *et al.*, 1984; Yap *et al.*, 1991; 1997; Lee *et al.*, 1993), until Vythilingam *et al.* (1997) found 2 females and 27 nymphs in a residence in Setapak Jaya in Kuala Lumpur and Lee & Lee (2000) trapped 2 specimens in a suburban apartment in Penang. In a study conducted in the animal facility of the Medical Faculty, University of Malaya, Sulaiman *et al.* (1993) trapped 1 female in the staff-resting area and 1 nymph in the food storage area, of this species. In a later study in the same faculty but conducted in the store and kitchen of the Clinical Student's Hostel, Jeffery *et al.* (2003b) trapped 878 *B. germanica*, followed by *P. americana* (n = 113) and *P. brunnea* (n = 12).

In peninsular Malaysia, *P. americana* is the largest domiciliary species. It is widely distributed but is rarely found or absent from Orang Asli (local aborigines) villages (Oothuman *et al.*, 1984; Sulaiman *et al.*, 1995; Jeffery *et al.*, 1999). *Periplaneta americana* was the dominant species in several surveys (Oothuman *et al.*, 1984; Yap *et al.*, 1991; Lee *et al.*, 1993; Lee & Lee, 2000). However, in the present and Jeffery *et al.* (2003) studies, it was superseded in dominance by *B. germanica*. *Periplaneta americana* has been seen entering in large numbers into human premises from the outside (unpublished observation). It has been recorded in large numbers in septic tanks in peninsular Malaysia (Anon, 1939 cited in Cornwell,

Table 1. Domiciliary cockroaches from restaurants in five zones of Kuala Lumpur Federal Territory, peninsular Malaysia

Species stages	Damansara (104)			Kepong (89)			Setapak (98)			Cheras (24)			Jalan Klang Lama (146)			Total (461)
	f	m	n	f	m	n	f	m	n	f	m	n	f	m	n	
<i>Blattella germanica</i>	673	972	2006	799	1033	969	651	821	1382	4	3	8	798	1088	1297	12504 [91.104%]
<i>Neostylopyga rhombifolia</i>	0	0	1	0	1	0	0	0	0	0	1	1	0	0	1	5 [0.036%]
<i>Periplaneta americana</i>	18	18	24	34	48	104	25	65	26	13	5	20	84	73	263	820 [5.974%]
<i>Periplaneta brunnea</i>	13	12	5	3	3	1	1	0	2	1	0	0	1	1	3	46 [0.335%]
<i>Supella longipalpa</i>	0	0	2	62	180	47	4	2	1	0	0	1	12	0	10	321 [2.339%]
<i>Symptloce pallens</i>	3	4	1	0	0	0	2	1	2	0	1	1	3	5	6	29 [0.211%]
	707	1006	2039	898	1265	1121	683	889	1413	18	10	31	898	1167	1580	13725

f- female; m-male; n-nymph; ( )-traps with cockroaches

1968). Predation of *P. americana* by the house shrew, *Suncus murinus* has been observed twice by one of us (JJ), and in one instance, the cockroach was completely consumed in less than 3 minutes. On another occasion, a toad, *Bufo melanostictus*, was observed at night in the same house. The next morning faeces believed to belong to the toad was seen and was collected and examined under a dissecting microscope. Large numbers of cockroach wings, estimated to belong to more than 10 individuals, was seen. The wings were considered to belong to *P. americana*, as this species was the only domiciliary species in that house at that time.

*Supella longipalpa*, a small sized domiciliary cockroach is generally found in drier areas of the household, particularly in bedroom, furnitures, television, etc. (Cornwell, 1968). Adults and nymphs have been trapped, on 3 occasions, from a booth of a car. Recently, a nymph was found in a cardboard box obtained from a provision shop.

*Periplaneta brunnea* in the past had been confused with *P. americana*, in peninsular Malaysia. Both these large *Periplaneta* spp. share superficial morphological similarities in the adult stages; however, are very distinct in the oothecal stage (Jeffery *et al.*, 1982). It is widespread but generally occurs in small numbers, except once, was the dominant species collected by Vythilingam *et al.* (1997), in Taman Ibu Kota and Setapak Jaya in Kuala Lumpur. *Periplaneta brunnea*, so far, has not been collected from aborigine villages (Oothuman *et al.*, 1984; Sulaiman *et al.*, 1995; Jeffery *et al.*, 1999).

*Symploce pallens*, a widely distributed species (Roth, 1984, 1996) was reported as a domiciliary pest species in peninsular Malaysia (Jeffery *et al.*, 1997; Lee *et al.*, 2000) and Singapore (Lee & Ng, 2009). In this study it was found in small numbers in all the 5 zones of the Kuala Lumpur Federal Territory. Vythilingam *et al.* (1997) collected it in

large numbers (identified as Blattellidae) in Taman Ibu Kota and Setapak Jaya, Kuala Lumpur. They isolated the following bacteria from this species: *Acinetobacter calcoaceticus*, *Enterobacter cloacae*, *Klebsiella p. pneumoniae* and *Klebsiella p. rhinocleromatis*.

Only 5 specimens of *N. rhombifolia* was collected from Kepong, Cheras and Jalan Kelang zones. Previously, it was found to occur together with *P. americana* and *P. brunnea* in the same niche (Jeffery *et al.*, 1982; Oothuman *et al.*, 1984; Yap *et al.*, 1991). Yap *et al.* (1991) hardly found them in rural areas. Jeffery *et al.* (2003a) collected them in small numbers in a rice-growing village in Kedah, where they were dominant over *Periplaneta australasiae*, *P. brunnea* and *S. longipalpa*. *Neostylopyga rhombifolia* was also collected from an aborigine resettlement in jungle at Pos Betau in Pahang state (Jeffery *et al.*, 1999). An important nosocomial pathogen, *Serratia marcescens* has been isolated from this species (Jeffery *et al.*, 1984).

Cockroaches have been reported as potential mechanical vectors of human diseases, shown to cause household allergy and entomophobia (Lee, 1997; Cochran, 1999). In this study 11 species of bacteria were isolated from *B. germanica*, *P. americana* and *P. brunnea* (Table 2).

In this study conducted in 5 zones of the Kuala Lumpur Federal Territory, 6 species of cockroaches were collected, mainly from the kitchen area of restaurants. The study areas were essentially urban with sparse vegetation. The species trapped were *B. germanica*, *N. rhombifolia*, *P. americana*, *P. brunnea*, *S. longipalpa* and *S. pallens*.

The following domiciliary species known from peninsular Malaysia, namely *Balta* (= *Lupparia*) *notulata*, *Nauphoeta cinerea*, *P. australasiae* and *Pycnoscelus surinamensis*, were not collected in this study.

Table 2. Bacteria isolated from domiciliary cockroaches collected from restaurants in Kuala Lumpur Federal Territory, peninsular Malaysia

Cockroach species	Bacteria species
<b><i>Blattella germanica</i></b>	
Specimen no. 1	<i>Enterobacter cloacae</i> <i>Klebsiella pneumoniae</i> ssp. <i>pneumoniae</i>
Specimen no. 2	<i>Enterobacter cloacae</i> <i>Klebsiella pneumoniae</i> ssp. <i>pneumoniae</i>
Specimen no. 3	<i>Enterobacter cloacae</i> <i>Klebsiella pneumoniae</i> ssp. <i>pneumoniae</i>
Specimen no. 4	<i>Enterobacter cloacae</i> <i>Klebsiella pneumoniae</i> ssp. <i>pneumoniae</i>
Specimen no. 5	<i>Serratia liquefaciens</i> <i>Klebsiella pneumoniae</i> spp. <i>rhinoscleromatis</i>
<b><i>Periplaneta brunnea</i></b>	
Specimen no. 1	<i>Acinetobacter calcoaceticus</i> var. <i>anitratus</i> <i>Klebsiella pneumoniae</i> ssp. <i>pneumoniae</i>
Specimen no. 2	<i>Escherichia vulneris</i>
Specimen no. 3	<i>Citrobacter diversus</i> / <i>amalonaticus</i>
<b><i>Periplaneta americana</i></b>	
Specimen no. 1	<i>Escherichia adecarboxylate</i> <i>Citrobacter freundii</i> <i>Proteus vulgaris</i>
Specimen no. 2	<i>Escherichia vulneris</i> <i>Klebsiella pneumoniae</i> ssp. <i>pneumoniae</i> <i>Klebsiella pneumoniae</i> ssp. <i>rhinoscleromatis</i> <i>Enterobacter agglomerans</i> 4

*Acknowledgement.* We gratefully acknowledge the support of our respective departments.

#### REFERENCES

- Anon. (1939). Cockroaches. Diseases to humans. *New Guinea Agriculture Gazette* **5**: 43-44 (Cited from Cornwell, 1968).
- Cochran, D.G. (1999). Cockroaches – Their biology, distribution and control. World Health Organization, Switzerland. *WHO/CDS/CPS/WHOPES/99.3*. pp.83.
- Cornwell, P.B. (1968). *The Cockroach*. Hutchinson and Company, London, pp.391.
- Cowan, S.T. & Steel, K.J. (1970). *Manual for Identification of Medical Bacteria*. London: Cambridge University Press, 238 pp.
- Jeffery, J., Oothuman, P., Zahedi, M.D. & Rampal, L. (1984). A simple cockroach trap. *Bulletin of the Public Health Society* **16**: 64-65.
- Jeffery, J., Nor Affandy, H., Sulaiman, A., Sallehudin, S., Norhayati, M., Baharudin, O., Hidayatulfathi, O., Ismail, M.G., Ahmad Firdaus, M.S. & Krishnasamy, M.

- (2003a). Domiciliary cockroaches and lizards and their parasites from Kg. Tebengau, a rice-growing village in Kedah. *Tropical Biomedicine* **20**(2): 199-201.
- Jeffery, J., Sulaiman, A., Khairul, A., Baharudin, O., Hidayatulfathi, O. & Krishnasamy, M. (2003b). Collection of domiciliary cockroaches and their parasites from the Faculty of Medicine, University of Malaya, Kuala Lumpur. *Tropical Biomedicine* **20**: 165-168.
- Jeffery, J., Vythilingam, I., Oothuman, P., Sulaiman, S., Zahedi, M., Eng, K.L. & Chiang, G.L. (1999). Domiciliary cockroaches from an Orang Asli (aborigine) resettlement in Pos Betau, Pahang State, Peninsular Malaysia. *Tropical Biomedicine* **16**: 35-38.
- Jeffery, J., Zahedi, M.D., Mohd Abdullah, M., Robiah, M. & Chiang, G.L. (1982). On the occurrence of the large brown cockroach *Periplaneta brunnea* in Peninsular Malaysia. *Malayan Nature Journal* **35**: 279-283.
- Jeffery, J., Zahedi, M., Oothuman, P., Sallehudin, S., Vythilingam, L. & Sulaiman, A. (1997). *Symploce pallens* (Stephens) (Dictyoptera: Blattellidae) – a new pest cockroach from Peninsular Malaysia. *Tropical Biomedicine* **14**: 137-139.
- Jeffery, J., Zahedi, M., Oothuman, P., Yusof, S., Busparani, V. & Robiah, M. (1984). Isolation of *Serratia marcescens* (Enterobacteriaceae) from cockroach, *Neostylopyga rhombifolia* (Blattaria). *Journal of Malaysian Society of Health* **4**: 57.
- Lee, C.Y. (1997). Medical importance of domiciliary cockroaches. *Singapore Microbiologist* **11**: 14-17.
- Lee, C.Y. (1998). Control of insecticide-resistant German cockroach, *Blattella germanica* (L.) (Dictyoptera: Blattellidae) in food-outlets with hydramethylmon-based bait stations. *Tropical Biomedicine* **15**: 45-51.
- Lee, C.Y. & Lee, L.C. (2000). Diversity of cockroach species and effect of sanitation level of cockroach infestation in residential premises. *Tropical Biomedicine* **17**: 39-43.
- Lee, C.Y., Chong, N.L. & Yap, H.H. (1993). A study on domiciliary cockroach infestation in Penang, Malaysia. *Journal of Bioscience* **2**: 95-98.
- Lee, C.Y., Khalid, M.S., Lee, L.C. & Chong, N.L. (2000). Notes on biological parameters of a new domiciliary pest, the smooth cockroach *Symploce pallens* (Stephens) (Dictyoptera: Blattellidae). *Tropical Biomedicine* **17**: 49-52.
- Lee, C.Y. & Ng, L.C. (2009). *Pest cockroaches of Singapore – A scientific guide for pest management professionals*. Singapore Pest Management Association. pp 64.
- Oothuman, P., Jeffery, J., Abdul Hamid, A.A., Edariah, A.B. & Jegathesan, M. (1989). Bacterial pathogens isolated from cockroaches trapped from paediatric wards in peninsular Malaysia. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **83**: 133-135.
- Oothuman, P., Jeffery, J., Zahedi, M.D., Rampal, L. & Shekhar, C. (1984). Distribution of different species of cockroaches in the district of Kelang, Selangor. *Journal of Malaysian Society of Health* **4**: 52-56.
- Roth, L.M. (1984). The genus *Symploce* Hebard. I. Species from the West Indies. (Dictyoptera: Blattaria, Blattellidae). *Entomologico Scandinavica* **15**: 25-63.
- Roth, L.M. (1996). Cockroaches from the Seychelles Island (Dictyoptera: Blattaria). *Journal of African Zoology* **110**: 97-128.
- Sulaiman, A., Jeffery, J., Oothuman, P., Shekhar, C. & Zahedi, D. (1995). Cockroaches and house lizard trapped from Malay and Orang Asli villages in Endau, Johore, Malaysia. *Tropical Biomedicine* **12**: 177-178.

- Sulaiman, A., Oothuman, P., Jeffery, J., Zahedi, M. & Baskaran, G. (1993). Report on cockroaches collected from an animal facility in Kuala Lumpur *Tropical Biomedicine* **10**: 197-198.
- Sulaiman, S., Cheon, Y.K., Aziz, A.H. & Jeffery, J. (2003). Isolations of bacteria pathogens from cockroaches trapped in downtown Kuala Lumpur. *Tropical Biomedicine* **20**: 53-57.
- Vythilingam, I., Jeffery, J., Oothuman, P., Abdul Razak, A.R. & Sulaiman, A. (1997). Cockroaches from urban human dwellings: isolation pathogens and control. *Southeast Asian Journal of Tropical Medicine and Public Health* **28**: 218-222.
- Yap, H.H., Chong, N.L., Loh, P.Y., Rohaizat, B. & Abdul Malik, Y. (1991). Survey of domiciliary cockroaches in Penang, Malaysia. *Journal of Bioscience* **2**: 71-75.
- Yap, H.H., Ong, C.H., Chong, N.L., Yahaya, A.M., Rahim, A.R., Awang, A.H. & Samsuri, O. (1997). Cockroach infestation in different household settlements in rural, suburban and urban areas on Penang Island, Malaysia. *Journal of Bioscience* **8**: 182-186.